

## Handout 05: Two-Samples

Consider extending our current setup for a statistical sample to the case where we have two samples, where each random observation is mutually independent from all the others:

$$X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{G}_X, \quad \mathbb{E}[X_i] = \mu_X, \quad \text{Var}[X_i] = \sigma_X^2$$

$$Y_1, \dots, Y_m \stackrel{iid}{\sim} \mathcal{G}_Y, \quad \mathbb{E}[Y_i] = \mu_Y, \quad \text{Var}[Y_i] = \sigma_Y^2$$

We will have the normal definitions of the sample means ( $\bar{X}$  and  $\bar{Y}$ ) and the sample variances ( $S_X^2$  and  $S_Y^2$ ). The **pooled variance**  $S_p^2$  is defined as a combination of the sample variance of  $X$  and  $Y$ :

$$S_p^2 = \frac{(n-1)S_X^2 + (m-1)S_Y^2}{n+m-2}$$

We will study its properties on today's worksheet. One of our key questions will how to build a point estimator and confidence interval for the difference in the means of the two distributions:  $\mu_X - \mu_Y$ .

### F-distribution

There is one more special distribution that we make significant usage of in statistics that, like the T-distribution, is particularly designed to serve as a pivot. If we have two independent random variables  $C_1$  and  $C_2$  that have chi-squared distributions of  $k_1$  and  $k_2$  degrees of freedom. Then, the following:

$$F = \frac{C_1/k_1}{C_2/k_2}$$

Has an F-distribution with  $k_1$  and  $k_2$  degrees of freedom (yes, it has two different degrees of freedom). We will use this today and then for several seemingly quite different tasks throughout the next several weeks.